

Global sections of structure sheafs of Keigher rings*

D. V. Trushin

Abstract

In answer to a question of J. Kovacic, we show that for any Keigher ring its differential spectrum coincides with differential spectrum of global sections of structure sheaf.

Introduction

The notion of structure sheaf $\mathcal{O}$ on differential spectrum X of differential ring A appeared in [2] and [3]. First disagreeable effects met is that global sections of structure sheaf does not necessarily coincide with initial ring. The question immediately arisen is: whether differential spectrum of global sections $\mathcal{O}(X)$ coincides with differential spectrum of initial ring A . Kovacic found conditions guaranteing that the desired equality holds [2, prop. 10.5]. However, the most interesting case is the case of Ritt algebras, because Ritt algebras appear in differential algebraic geometry. Example [3, ex. 7.4] shows that proposition [2, 10.5] cannot be applied to all Ritt algebras. We give the answer for more wide class of differential rings – the class of Keigher rings.

The idea of the proof is following. We change structure sheaf to other one which is not so hard for investigation. After this changing we do not lose the information about differential spectrum. Using simple topological technic, we show that differential spectrum of global sections of new sheaf coincides with that of initial ring. In our method the condition that the ring is a Keigher ring is essentially used.

In the first section we recall some basic definitions. All terms are taken from [2] and [3]. In the next section we construct auxiliary sheaf and investigate its properties. Topological technic is presented in lemma 1 and the main result of the section is statement 6. In the last section we prove main result – theorem 8. Corollary 9 gives the answer in the case of Ritt algebras.

*This paper is dedicated to Jerald J. Kovacic

1 Terms and notation

Throughout the text the word ring means an associative commutative ring with an identity. All homomorphisms preserve the identity element. A differential ring is a ring with finitely many pairwise commuting derivations. A ring is a Keigher ring if for any differential ideal $\mathfrak{a}$ its radical $\mathfrak{r}(\mathfrak{a}) = \{x \mid x^n \in \mathfrak{a}\}$ is a differential ideal too. For any differential ring A the set of all prime differential ideals is denoted by $\text{Spec}^\Delta A$ and is called a differential spectrum. For any subset $E \subseteq A$ the set of all prime differential ideal containing E is denoted by $V(E)$. Defining $V(E)$ as closed subsets, we provide differential spectrum with topology.

Denote differential spectrum of A by X and let s be an element of A . Then the set of all prime differential ideals not containing s is denoted by X_s . For the set $S = \{s^n\}_{n=0}^\infty$ the localization $S^{-1}A$ is denoted by A_s . If $\mathfrak{p}$ is a prime ideal then by $A_{\mathfrak{p}}$ we denote the ring $S^{-1}A$, where $S = A \setminus \mathfrak{p}$. For any differential homomorphism $f: A \rightarrow B$ we define the corresponding mapping $f^*: \text{Spec}^\Delta B \rightarrow \text{Spec}^\Delta A$ by the rule $f^*(\mathfrak{p}) = f^{-1}(\mathfrak{p})$.

We shall need the notion of structure sheaf on differential spectrum. We shall briefly recall its definition. Consider the set of all functions

$$\{f: X \rightarrow \bigcup_{\mathfrak{p} \in X} A_{\mathfrak{p}} \mid f(\mathfrak{p}) \in A_{\mathfrak{p}}\}.$$

Function f will be called a regular at the point $\mathfrak{p}$ if there exist an open neighborhood U of $\mathfrak{p}$ and elements $a, b \in A$, where for all $\mathfrak{q} \in U$ we have $b \notin \mathfrak{q}$, such that for all $\mathfrak{q} \in U$ it follows that $f(\mathfrak{q}) = a/b$ in $A_{\mathfrak{q}}$. The set of all functions that regular at all points of open subset U will be denoted by $\mathcal{O}(U)$. The family $\mathcal{O}(U)$ with restriction homomorphisms form a sheaf of differential rings. We shall call this sheaf a structure sheaf.

The ring $\mathcal{O}(X)$ will be denoted by $\hat{A}$. There is a mapping $\iota: A \rightarrow \hat{A}$ such that $a \mapsto \varphi$, where $\varphi(\mathfrak{p}) = a/1$ in $A_{\mathfrak{p}}$. It is easy to see that ι is a differential homomorphism.

All details about structure sheaf can be found in [2]. We assume that the reader is acquainted with this work.

2 Auxiliary sheaf

In this section we construct another sheaf for which the desired problem will be solved by straightforward calculation. The general proof based on the reduction of the initial problem to the problem solved in this section.

Let A be an arbitrary differential ring. For any prime differential ideal $\mathfrak{p}$ its residue field will be denoted by $K(\mathfrak{p})$, i. e., the fraction field of $A/\mathfrak{p}$. Consider the set of all functions

$$\{f: X \rightarrow \bigcup_{\mathfrak{p} \in X} K(\mathfrak{p}) \mid f(\mathfrak{p}) \in K(\mathfrak{p})\}.$$

The function will be called regular at the point $\mathfrak{p}$ if there exist an open neighborhood U of $\mathfrak{p}$ and elements $a, b \in A$, where for all $\mathfrak{q} \in U$ we have $b \notin \mathfrak{q}$, such that for all $\mathfrak{q} \in U$ it follows that $f(\mathfrak{q}) = a/b$ in $K(\mathfrak{q})$. The set of all functions that regular at all points of U will be denoted by $\mathcal{O}'(U)$. The family of all such rings with restriction homomorphisms form a sheaf of differential rings on X .

Define a differential homomorphism $\iota': A \rightarrow \mathcal{O}'(X)$ by the rule $a \mapsto \varphi$ such that $\varphi(\mathfrak{p}) = a$ in $K(\mathfrak{p})$. Our proof based on the following simple statement.

Lemma 1. *Let $\varphi: X \rightarrow Y$ be a mapping between topological spaces and let two covers U_α and V_α of X and Y respectively be given. Then if for any pair of indexes α and α' the mapping $\varphi: U_\alpha \cap U_{\alpha'} \rightarrow V_\alpha \cap V_{\alpha'}$ is well-defined and is a homeomorphism, then $\varphi: X \rightarrow Y$ is a homeomorphism too.*

Proof. Let us take $\alpha = \alpha'$, then φ is a local homeomorphism. We need to prove that φ is injective. Let $x \in U_\alpha$ and $y \in U_{\alpha'}$ be such that $z = f(x) = f(y)$. Then $z \in V_\alpha \cap V_{\alpha'}$. Since φ is a homeomorphism between $U_\alpha \cap U_{\alpha'}$ and $V_\alpha \cap V_{\alpha'}$, then the intersection $U_\alpha \cap U_{\alpha'}$ is not empty and contains w such that $\varphi(w) = z$. But x and w both belong to U_α and their images coincide, thus $x = w$. The same reasons show that $y = w$. $\square$

Now we shall prove some auxiliary facts.

Lemma 2. *Let A and B be differential rings, $\nu: A \rightarrow B$ be a differential homomorphism, and B is differentially generated over A by one single element η such that there exists a family of elements $b_1, \dots, b_n$ of A with conditions $b_k \eta \in \nu(A)$ and $\{b_1, \dots, b_n\} = A$. Then the contraction mapping $\nu^*: \text{Spec}^\Delta B \rightarrow V(\ker \nu)$ is a homeomorphism.*

Proof. First of all, changing A onto $A/\ker \nu$, we may suppose that ν is injective. Now we shall divide differential spectra to finitely many open subsets as follows

$$\begin{aligned} X &= X_{b_1} \cup \dots \cup X_{b_n} \\ Y &= Y_{b_1} \cup \dots \cup Y_{b_n} \end{aligned}$$

We need to check the condition of previous lemma for coverings X_{b_i} and Y_{b_i} . Proof is based on [1, ex. 21 (i,ii) p. 46]. We shall replace X_{b_i} by differential spectrum of A_{b_i} , Y_{b_i} by differential spectrum of B_{b_i} , and from the mentioned exercise it follows that restriction of ν^* coincides with $\nu_{b_i}^*$ (see [1, ex. 21 (ii) p. 47]).

In order to show the promised fact we shall calculate both localizations A_{b_i} and B_{b_i} . Let $b_i \eta = a_i \in A$. Consider

$$B_{b_i} = (A\{\eta\})_{b_i} = A\{a_i/b_i, 1/b_i\} = A\{1/b_i\} = A_{b_i}$$

Therefore localization of $\nu: A \rightarrow B$ by b_i induces the identity mapping

$$\nu_{b_i} = Id: A_{b_i} \rightarrow B_{b_i} = A_{b_i}.$$

So, the desired homeomorphism between X_{b_i} and Y_{b_i} has been obtained.

The last step of the proof is to show that ν^* induces a homeomorphism between $X_{b_i} \cap X_{b_j}$ and $Y_{b_i} \cap Y_{b_j}$. Having noted that $X_{b_i} \cap X_{b_j} = X_{b_i b_j}$ and $Y_{b_i} \cap Y_{b_j} = Y_{b_i b_j}$ and applying previous arguments with $b_i b_j$ instead of b_i , we get the result. $\square$

Using induction we show the following.

Lemma 3. *Let A and B be differential rings, $\nu: A \rightarrow B$ be a differential homomorphism, and B be differentially finitely generated over A such that for any generator η there exists a family of elements $b_1, \dots, b_n$ with conditions $b_k \eta \in \nu(A)$ and $\{b_1, \dots, b_n\} = A$. Then $\nu^*: \text{Spec}^\Delta B \rightarrow V(\ker \nu)$ is a homeomorphism.*

Proof. Let B be generated over A by the elements $\eta_1, \dots, \eta_n$, and let B_1 be subring of B generated over A by elements $\eta_1, \dots, \eta_{n-1}$. Then homomorphism ν can be presented as the following composition

$$A \xrightarrow{\nu_1} B_1 \xrightarrow{\nu_2} B.$$

From the previous lemma $V(\ker \nu_2)$ is homeomorphic to $\text{Spec}^\Delta B$. From inductive assumption $\text{Spec}^\Delta B_1$ and $V(\ker \nu_1)$ are homeomorphic, and thus their corresponding subspaces $V(\ker \nu_2)$ and $V(\ker \nu)$ are homeomorphic too. Consequently, $V(\ker \nu)$ and $\text{Spec}^\Delta B$ are homeomorphic to each other. $\square$

Lemma 4. *Let A_α be direct system of differential A -algebras and $\nu_\alpha: A \rightarrow A_\alpha$ be corresponding homomorphisms. Suppose that for any α the mapping ν_α^* is a homeomorphism. Then ν^* is a homeomorphism, where $\nu: A \rightarrow \varinjlim_\alpha A_\alpha$ is the direct limit of ν_α .*

Proof. Let $\varinjlim_\alpha A_\alpha$ be denoted by A' . Firstly we show that ν is bijective. Surjectivity: consider a prime differential ideal $\mathfrak{p}$ in A . Let $\mathfrak{p}_\alpha$ be corresponding to $\mathfrak{p}$ ideals in A_α . It is clear that the ideal $\varinjlim_\alpha \mathfrak{p}_\alpha$ is a prime differential ideal in A' contracting to $\mathfrak{p}$.

Injectivity: suppose that there exist two prime differential ideals $\mathfrak{q}$ and $\mathfrak{q}'$ in A' contracting to $\mathfrak{p}$. Then there is an element $s \in A'$ such that $s \in \mathfrak{q}' \setminus \mathfrak{q}$. From [1, chapter 2, ex. 14] it follows that there are α and element s_α in A_α such that s_α maps to s . Let $\mathfrak{q}_\alpha$ and $\mathfrak{q}'_\alpha$ denote contractions of $\mathfrak{q}$ and $\mathfrak{q}'$ on A_α respectively. Then we have $s_\alpha \in \mathfrak{q}'_\alpha \setminus \mathfrak{q}_\alpha$. But both these ideals contract to $\mathfrak{p}$ and belong to A_α , contradiction.

Now we shall show that ν is a homeomorphism. We just need to prove that the image of any principal open set is open. Let s be an arbitrary element of A' , we shall show that the image of $\text{Spec}^\Delta(A')_s$ is open in $\text{Spec}^\Delta A$. Indeed, for some α there is an element s_α in A_α such that its image coincides with s . Then $\text{Spec}^\Delta(A_\alpha)_{s_\alpha} = \text{Spec}^\Delta(A')_s$. But by the data the image of $\text{Spec}^\Delta(A_\alpha)_{s_\alpha}$ is open in $\text{Spec}^\Delta A$, q. e. d. $\square$

Lemma 5. *For every element $\eta \in \mathcal{O}'(X)$ the following holds. There exists a family of elements $b_1, \dots, b_m \in A$ such that $b_i \eta \in \iota'(A)$ and $\{b_1, \dots, b_m\} = A$.*

Proof. Indeed, let $U_1 \cup \dots \cup U_m = X$ be a covering of differential spectrum such that for any $\mathfrak{p} \in U_i$ we have $\eta(\mathfrak{p}) = a_i/b_i$ in $K(\mathfrak{p})$. We shall change a_i onto $a_i b_i$ and b_i^2 . Then for any $\mathfrak{p} \notin U_i$ we have $a_i = 0$ in $K(\mathfrak{p})$. Without loss of generality we may suppose that $U_i = X_{b_i}$. Then, since all U_i cover X it follows that $\{b_1, \dots, b_m\} = A$. Let us show that for every $\mathfrak{p}$ the condition $b_i \eta(\mathfrak{p}) = a_i$ in $K(\mathfrak{p})$ holds. Indeed, if $\mathfrak{p} \in U_i$, then $b_i \eta(\mathfrak{p}) = b_i(a_i/b_i) = a_i$ in $K(\mathfrak{p})$. If $\mathfrak{p} \in U_j \setminus U_i$, then $b_i \eta(\mathfrak{p}) = b_i(a_j/b_j) = 0$ in $K(\mathfrak{p})$, but $a_i = 0$ in $K(\mathfrak{p})$ too. Consequently, $b_i \eta = \iota'(a_i)$. $\square$

Statement 6. *Let D be a differential subring of $\mathcal{O}'(X)$ containing image of A , then we are able to consider the homomorphism $\iota': A \rightarrow D$. In that case the following mapping is a homeomorphism.*

$$\iota'^*: \text{Spec}^\Delta D \rightarrow \text{Spec}^\Delta A.$$

Proof. From the previous lemma it follows that for any differentially finitely generated over A subalgebra in $\mathcal{O}'(X)$ the conditions of lemma 3 holds. From the definition of ι' it follows that $\ker \iota'$ belongs to the intersection of all prime differential ideals of A . Consequently, for any differentially finitely generated algebra D the statement is proven. But any differential algebra can be presented as a direct limit of its differentially finitely generated subalgebras. Then from lemma 4 it follows that statement holds for any subalgebra D . $\square$

3 Main result

In this section we reduce the main result to statement 6 proven. We have two sheafs on topological space X , they are $\mathcal{O}$ and $\mathcal{O}'$. Consider the homomorphism of sheafs $\varphi(U): \mathcal{O}(U) \rightarrow \mathcal{O}'(U)$ defined by the rule $\varphi(f)(\mathfrak{p}) = f(\mathfrak{p}) \bmod \mathfrak{p}$. From now we suppose that the given ring A is a Keigher ring.

Statement 7. *With the above conventions, we have that $\ker \varphi(X)$ coincides with the nilradical of $\mathcal{O}(X)$.*

Proof. By the definition $\ker \varphi(X)$ coincides with

$$\{f \in \mathcal{O}(X) \mid f(\mathfrak{p}) \in \mathfrak{p}A_{\mathfrak{p}}\}.$$

Let us show that nilradical lays in kernel. Let f be a nilpotent element, then for any $\mathfrak{p}$ we have $f^n(\mathfrak{p}) = 0$. So, $f^n(\mathfrak{p}) \in \mathfrak{p}A_{\mathfrak{p}}$. Consequently, $f(\mathfrak{p}) \in \mathfrak{p}A_{\mathfrak{p}}$.

Conversely, let $f \in \ker \varphi(X)$, then there exists a covering $U_1 \cup \dots \cup U_n = X$ such that for every i we have $f(\mathfrak{p}) = a_i/b_i$ in $A_{\mathfrak{p}}$, whenever $\mathfrak{p} \in U_i$. Without loss of generality we may suppose that $U_i = X_{b_i}$. Let us fix some i , then $f(\mathfrak{p}) = a_i/b_i \in \mathfrak{p}A_{\mathfrak{p}}$ for all $\mathfrak{p}$ not containing b_i . Then $a_i \in \mathfrak{p}$ for all $\mathfrak{p}$ not containing b_i . Therefore, a_i belongs to the intersection of all prime ideals in A_{b_i} .

Since A is a Keigher ring (A_{b_i} too) a_i belongs to nilradical of A_{b_i} . Hence, for some k_i we have $a_i^{k_i} = 0$ in A_{b_i} . Therefore, for some m_i it follows that $b_i^{m_i} a_i^{k_i} = 0$ in A . And so, we have that $(a_i b_i)^{n_i} = 0$ for some n_i . Let $n = \max_i n_i$, then for every i we have $(a_i b_i)^n = 0$ in A . Consequently, for every i and every $\mathfrak{p} \in U_i$ the following holds

$$f^n(\mathfrak{p}) = \left(\frac{a_i}{b_i} \right)^n = \frac{(a_i b_i)^n}{b_i^{2n}} = 0 \text{ in } A_{\mathfrak{p}}.$$

□

Theorem 8. *Let D be subring of $\mathcal{O}(X)$ containing image of A , then we are able to consider homomorphism $\iota: A \rightarrow D$. In that case the following mapping is a homeomorphism*

$$\iota^*: \text{Spec}^\Delta D \rightarrow \text{Spec}^\Delta A.$$

Proof. Consider homomorphism $\varphi(X): D \rightarrow \mathcal{O}'(X)$, from the previous statement it follows that the kernel of $\varphi(X)$ coincides with the nilradical of D . Denote the quotient of D by its nilradical by D' . It is clear that $\text{Spec}^\Delta D$ can be identified with $\text{Spec}^\Delta D'$. We have the following sequence of homomorphisms

$$A \rightarrow D' \rightarrow \mathcal{O}'(X).$$

And the composition coincides with ι' . From statement 6 it follows that $\text{Spec}^\Delta D' \rightarrow \text{Spec}^\Delta A$ is a homeomorphism. Remembering that $\text{Spec}^\Delta D$ is homeomorphic to $\text{Spec}^\Delta D'$, we get the desired result. □

Corollary 9. *Let A be a Ritt algebra, and let $\iota: A \rightarrow \hat{A}$ be the canonical homomorphism. Then the mapping*

$$\iota^*: \text{Spec}^\Delta \hat{A} \rightarrow \text{Spec}^\Delta A$$

is a homeomorphism.

References

- [1] M. F. Atiyah, I. G. Macdonald. Introduction to commutative algebra. Addison-Wesley. 1969.
- [2] J. J. Kovacic. Differential Schemes, Differential Algebra and Related Topics, Newark, NJ, 2000, pp. 71–94.
- [3] J. J. Kovacic. Global sections of diffspec, J. Pure and Applied Alg., 171 (2002), 265–288